

Coding Spotlight: Hypertension -- A providers' guide for coding

Published: Jan 1, 2020 - State & Federal / Medicaid

Category: Medicaid

ICD-10-CM hypertension coding highlights:

- Hypertensive crisis can involve hypertensive urgency or emergency.
- Hypertension can occur with heart disease, chronic kidney disease (CKD) or both.
- ICD-10-CM classifies hypertension by type as essential or primary (categories I10-I13) and secondary (category I15).¹
- Categories I10-I13 classify primary hypertension according to a hierarchy of the disease from its vascular origin (I10) to the involvement of the heart (I11), CKD (I12), or heart and CKD combined (I13).¹

Hypertension categories:

Code	Description
I10	Essential (primary) hypertension
I11.0	Hypertensive heart disease with heart failure
I11.9	Hypertensive heart disease without heart failure
I12.0	Hypertensive CKD with stage 5 CKD or end-stage renal disease (ERSD)
I12.9	Hypertensive CKD with stage 1 through stage 4 CKD or unspecified CKD
I13.0	Hypertensive heart and CKD with heart failure and stage 1 through stage 4 CKD or unspecified CKD
I13.10	Hypertensive heart and CKD without heart failure with stage 1 through stage 4 CKD or unspecified CKD
I13.11	Hypertensive heart and CKD without heart failure with stage 5 CKD or ERSD
I13.2	Hypertensive heart and CKD with heart failure and with stage 5 CKD or ERSD
I15.-	Secondary hypertension
I16.-	Hypertensive crisis

Hypertensive heart disease

ICD-10-CM presumes a causal relationship between hypertension and heart involvement and classifies hypertension and heart conditions to category I11 (hypertensive heart disease) because the two conditions are linked by the term “with” in the *Alphabetic Index of ICD-10-CM*. These conditions should be coded as related even in the absence of provider documentation linking them. Code first I11.0 (hypertensive heart disease with heart failure) as instructed by the note at category I50 (heart failure). If the provider specifically documents different causes for the hypertension and the heart condition, the heart condition (I50.-, I51.4 to I51.x9) and hypertension are coded separately.¹

Category I11 is subdivided to indicate whether heart failure is present. However, an additional code from category I50 is required to specify the type of heart failure, if known.

Documentation may vary, but coding instructions remain the same. For example:

- Congestive heart failure due to hypertension: I11.0 + I50.9
- Hypertensive heart disease with congestive heart failure: I11.0 + I50.9
- Congestive heart failure with hypertension: I11.0 + I50.9

Other heart conditions that have an assumed causal connection to hypertensive heart disease:

Code	Description
I51.4	Myocarditis, unspecified
I51.5	Myocardial degeneration
I51.7	Cardiomegaly
I51.81	Takotsubo syndrome
I51.89	Other ill-defined heart diseases
I51.9	Heart disease, unspecified

Hypertension and CKD

When the diagnostic statement includes both hypertension and CKD, ICD-10-CM assumes there is a cause-and-effect relationship. A code from category I12 (hypertensive CKD) is assigned because the two conditions are linked by the term “with” in the *Alphabetic Index of ICD-10-CM*. These conditions should be coded as related even in the absence of provider

documentation linking them, unless the documentation clearly states the conditions are unrelated.¹

A fourth character is used with category I12 to indicate the stage of the CKD. The appropriate code from category N18 should be used as a secondary code to identify the stage of CKD.

Hypertensive heart and CKD

Combination category I13 codes are assigned for hypertensive heart and CKD when there is hypertension with both heart and kidney involvement. If heart failure is present, an additional code from category I50 is assigned to identify the type of heart failure.¹

The appropriate code from category N18 (CKD) should be used as secondary code with a code from category I13 to identify the stage of CKD.

Hypertensive cerebrovascular disease

For hypertensive cerebrovascular disease, first the appropriate code from categories I60 to I69 is assigned followed by the hypertension code.

Hypertensive retinopathy

Subcategory H35.0 (background retinopathy and retinal vascular changes) should be used with a code from category I10 to I15 (hypertensive disease to include the systemic hypertension).²

Hypertension, secondary

Two codes are required -- one to identify the underlying etiology and one from category I15 to identify the hypertension. For example:

- Hypertension due to systemic lupus erythematosus: M32.10 + I15.8
- Acromegaly with secondary hypertension seen for hypertension management: I15.2 + E22.0

Hypertension, transient

Code R03.0 (elevated blood pressure reading without diagnosis of hypertension) is assigned unless the patient has an established diagnosis of hypertension. For transient hypertension of pregnancy, code O13.- (gestational [pregnancy-induced] hypertension without significant proteinuria) or O14.- (pre-eclampsia).

Hypertensive crisis

A code from category I16 (hypertensive crisis) is assigned for any documented hypertensive urgency, hypertensive emergency or unspecified hypertensive crisis. Report two codes, at a minimum, for hypertensive crisis. The crisis code is reported in addition to the underlying hypertension code (I10 to I15).¹

- Hypertensive urgency: I16.0
- Hypertensive emergency: I16.1
- Hypertensive crisis, unspecified: I16.9

Pulmonary hypertension

Pulmonary hypertension is classified to category I27 (other pulmonary heart diseases). For secondary pulmonary hypertension (I27.1, I27.2-), any associated conditions or adverse effect of drugs or toxins should be coded.²

More coding tips

Blood pressure and medication management should be assessed at every encounter involving a hypertensive patient. Clarity is important in documenting hypertension. Ensure that the diagnosis is captured by noting it in the medical record documentation:

- Specify a pregnant patient with hypertension as having a pre-existing, gestational, pre-eclampsic or eclampsic hypertension.
- Document and code the smoking status of a patient with hypertension:
 - Current smoker: F17.
 - Personal history of tobacco dependence: Z87.891
 - Tobacco use: Z72.0
 - Exposure to environmental tobacco smoke: Z57.31
- Document any causal relationship between hypertension and background retinopathy or other condition in which the hypertension caused vascular changes and organ damage.

HEDIS Quality Measures for hypertension

The Controlling High Blood Pressure (CBP) measure looks at a sample of members ages 18 to 85 years of age who have a diagnosis of hypertension and whose blood pressure (BP) is regularly monitored and controlled.³

HEDIS is a registered trademark of the National Committee for Quality Assurance (NCQA).

Record your efforts

Document blood pressure and diagnosis of hypertension. Patients whose BP is adequately controlled include patients ages 18 to 59 with less than 140/90 mm Hg.

Both systolic and diastolic values must be below the stated value. The most recent BP measurement during the year counts toward compliance.

What does not count?

- A BP measurement taken on the same day or one day before the test or procedure (fasting blood tests not included).
- Patient reported BP measurements.
- A BP measurement taken on the same day as a diagnostic test or procedure that requires a change in diet or medication regimen. For example:
 - Procedures that require a change in diet or medication regimen: colonoscopy, dialysis, infusions, chemotherapy, nebulizer treatment with albuterol and injection of lidocaine prior to mole removal
 - Procedures (low-intensity or preventive) that would not disqualify the BP reading: vaccinations, injections, TB test, intrauterine device insertion and eye exam with dilating agents

Codes to identify hypertension

ICD-10-CM	CPT Category II codes ⁴
I10	3074F: systolic BP <130
	3075F: systolic BP 130 to 139
	3077F: systolic BP ≥140
	3078F: diastolic BP <80
	3079F: diastolic BP 80 to 89

Strategies for success

- Improve the accuracy of BP measurements performed by your clinical staff by:
 - Providing training materials from the American Heart Association.
 - Conducting BP competency tests to validate the education of each clinical staff member.
 - Making a variety of cuff sizes available.
- Instruct your office staff to recheck BPs for all patients with initial recorded readings greater than systolic 140 mm Hg and diastolic of 90 mm Hg during outpatient office visits; have your staff record the recheck in the patient's medical records.
- Educate your patients (and their spouses, caregivers or guardians) about the elements of a healthy lifestyle, such as:
 - Heart-healthy eating and low-salt diet.
 - Smoking cessation and avoiding secondhand smoke.
 - Adding regular exercise to daily activities.
 - Home BP monitoring.
 - Ideal body mass index.
 - The importance of taking all prescribed medications as directed.
- Remember to include the applicable Category II reporting codes on the claim form to help reduce the burden of HEDIS medical record review.

Resources

"ICD-10-CM Expert for Physicians. The complete official code set," Optum360, LLC (2019).

Elsevier, "ICD-10-CM/PCS Coding, Theory and Practice — 2019/2020 Edition."

"HEDIS Measures and Technical Resources," NCQA, accessed April 15, 2019,

<https://www.ncqa.org/hedis/measures>.

"CPT 2019 Professional Edition," American Medical Association (2019).

"HCPCS Level II," American Medical Association (2019).

ANV-NU-0055-19 October 2019

URL: <https://providernews.anthem.com/nevada/article/coding-spotlight-hypertension-a-providers-guide-for-coding-1>

Featured In:

January 2020 Anthem Provider News and Important Updates - Nevada

Anthem Blue Cross and Blue Shield is the trade name of: In Colorado Rocky Mountain Hospital and Medical Service, Inc. HMO products underwritten by HMO Colorado, Inc. In Connecticut: Anthem Health Plans, Inc. In Georgia: Blue Cross Blue Shield Healthcare Plan of Georgia, Inc. In Indiana: Anthem Insurance Companies, Inc. In Kentucky: Anthem Health Plans of Kentucky, Inc. In Maine: Anthem Health Plans of Maine, Inc. In Missouri (excluding 30 counties in the Kansas City area): RightCHOICE® Managed Care, Inc. (RIT), Healthy Alliance® Life Insurance Company (HALIC), and HMO Missouri, Inc. RIT and certain affiliates administer non-HMO benefits underwritten by HALIC and HMO benefits underwritten by HMO Missouri, Inc. RIT and certain affiliates only provide administrative services for self-funded plans and do not underwrite benefits. In Nevada: Rocky Mountain Hospital and Medical Service, Inc. HMO products underwritten by HMO Colorado, Inc. dba HMO Nevada. In New Hampshire: Anthem Health Plans of New Hampshire, Inc. HMO plans are administered by Anthem Health Plans of New Hampshire, Inc. and underwritten by Matthew Thornton Health Plan, Inc. In Ohio: Community Insurance Company. In Virginia: Anthem Health Plans of Virginia, Inc. trades as Anthem Blue Cross and Blue Shield in Virginia, and its service area is all of Virginia except for the City of Fairfax, the Town of Vienna, and the area east of State Route 123. In Wisconsin: Blue Cross Blue Shield of Wisconsin (BCBSWI), which underwrites or administers the PPO and indemnity policies and underwrites the out of network benefits in POS policies offered by CompCare or WCIC; CompCare Health Services Insurance Corporation (CompCare) underwrites or administers the HMO policies and Wisconsin Collaborative Insurance Company (WCIC) underwrites or administers Well Priority HMO or POS policies. Independent licensees of the Blue Cross and Blue Shield Association. Anthem is a registered trademark of Anthem Insurance Companies, Inc. The Blue Cross and Blue Shield names and symbols are registered marks of the Blue Cross and Blue Shield Association. Use of the Anthem websites constitutes your agreement with our Terms of Use.
